

Strongly Semistable Bundles on a Curve over a finite field

S. Subramanian

Abstract

We show that a principal G bundle on a smooth projective curve over a finite field is strongly semistable if and only if it is defined by a representation of the fundamental group scheme of the curve into G .

MSC 2000 classification:14-XX

Keywords:semistable bundles,fundamental group scheme.

1. Introduction

Let G be a semisimple algebraic group and X a smooth projective curve, both defined over a finite field k of characteristic p . Let $E \rightarrow X$ be a principal G -bundle on X . E is said to be stable (resp. semistable) if for every reduction of structure group $E_P \subset E$ to a parabolic subgroup P of G , we have

$$\deg E_P(\mathfrak{p}) < (\text{resp. } \leq) \deg E_G(\mathfrak{g})$$

where $\mathfrak{g}$ and $\mathfrak{p}$ denote the lie algebras of G and P respectively, and $E_P(\mathfrak{p})$ (respectively $E_G(\mathfrak{g})$) denotes the Lie algebra bundle associated to E_P (respectively E_G) by the adjoint action of P (respectively G) on $\mathfrak{p}$ (respectively $\mathfrak{g}$). When the base

field is the field of complex numbers, the theorem of Narasimhan-Seshadri-Ramanathan says that a stable G -bundle is defined by a unitary representation of the topological fundamental group of X . In this article, we show that in the case when k is a finite field, a strongly semistable principal G -bundle on X (recall that a bundle is said to be strongly semistable if all of its Frobenius pull backs are semistable) is defined by a representation of the fundamental group scheme $\pi(X) \rightarrow G$. Here, by fundamental group scheme, we mean the proalgebraic group scheme introduced by M. Nori in [3], [4]. The author wishes to thank the referee for his helpful comments on an earlier version of this paper.

2. Preliminaries

Let k be a finite field of characteristic p and let X be a smooth projective curve defined over k . We assume that X has a rational point (say x) defined over k . A vector bundle $V \rightarrow X$ (over k) of degree zero is said to be essentially finite if there exists a finite group scheme H over k and a principal H -bundle $\pi : E_0 \rightarrow X$ on X , such that π^*V is trivial on E_0 (see Proposition(3.10),page 83,in [5]). Let $\mathcal{C}$ denote the category of all essentially finite vector bundles defined on X over k . We define a fibre functor T on $\mathcal{C}$ by defining $T(V) = V_x$, the fibre of V at x , for every $V \in \mathcal{C}$. With this fibre functor, $\mathcal{C}$ is a Tannaka category over k (see section 2.3 in [5]). The associated affine group scheme $\pi(X, x)$ is called the fundamental group scheme of X over k . Further, there is a principal $\pi(X, x)$ bundle $\mathcal{X} \rightarrow X$ over k (see section 2.3,page 84,in [5]). Given a semisimple algebraic group G over k , and an algebraic group

homomorphism $\pi(X, x) \rightarrow G$ (such a homomorphism factors through a finite group scheme quotient of $\pi(X, x)$), there is associated a principal G -bundle $E \rightarrow X$, associated to $\mathcal{X} \rightarrow X$ by the homomorphism $\pi(X, x) \rightarrow G$.

We now consider a principal G bundle $E \rightarrow X$, which we assume to be strongly semistable. Since G is assumed to be semisimple, we can find a G -module V such that $G \subset SL(V)$. The associated vector bundle $E(V)$ is strongly semistable by the theorem of Ramanan-Ramanathan (see Theorem 3.23 in [6]). We form the category $\mathcal{M}$ whose objects are subquotients of degree zero of finite direct sums

$$\bigoplus_i E(V)^{\oplus n_i} \oplus E(V)^{* \oplus m_i}$$

where n_i, m_i are nonnegative integers. The homomorphisms in $\mathcal{M}$ are all homomorphisms of the respective vector bundles on X . Given a vector bundle W in $\mathcal{M}$, we can associate to it its fibre W_x at x , and with this as the fibre functor, $\mathcal{M}$ is a Tannaka category over k (see Section 4 in [1]). There is an associated affine group scheme M , called the monodromy group scheme of E . Further, if $Vect(X)$ denotes the category of vector bundles on X , then there is a naturally defined functor $\phi : \mathcal{M} \rightarrow Vect(X)$ which associates to an object of $\mathcal{M}$ the vector bundle on X that it represents. The functor ϕ satisfies the axioms of Lemma(2.1), page 77, in [5]. Hence, by Proposition(2.5), page 78, in [5], we obtain a principal M -bundle $E_M \rightarrow X$ on X , defined over k (see [1], section 4). Further, since k is a finite field, $H^1(k, G)$ is trivial (see Theorem 1', section 2.3, Chapter 3, page 132 in [7]), and hence we can assume that there is a k rational point in the fibre E_x of E at x . With this choice of point in E_x , we obtain an embedding $M \subset G$, and a reduction of structure group $E_M \subset E$ (see section 4 in [1] for details). The monodromy group

scheme M and the bundle E_M do not depend on the choice of representation $G \subset SL(V)$, and the bundle E_M is characterised as the minimal degree zero reduction of structure group of E (see Proposition 4.9, Corollary 4.10 and Proposition 4.11 in [1]).

3. Main Theorem

Let k be a finite field and let X be a smooth projective curve over k . We assume that X has a rational point x over k . Let G be a semisimple algebraic group over k . Let F denote the Frobenius morphism on X .

Definition (3.1) A principal G bundle $E \rightarrow X$ on X over k is said to be semistable if, for every reduction of structure group $E_P \subset E$ to a parabolic subgroup P of G , we have

$$\text{degree } E_P(\mathfrak{p}) \leq 0.$$

The principal bundle is said to be strongly semistable if $F^{n*}E$ is semistable for all $n \geq 1$. We can now prove

Theorem (3.2) Let $E \rightarrow X$ be a strongly semistable G -bundle on X over k . Then $E \rightarrow X$ is defined by a homomorphism $\pi(X, x) \rightarrow G$. This homomorphism $\pi(X, x) \rightarrow G$ is unique upto inner conjugation by an element of $G(k)$. Further, if G' is another semisimple algebraic group over k and $G \rightarrow G'$ a homomorphism of algebraic groups, then the strongly semistable G' bundle $E_{G'} \rightarrow X$ associated to $E \rightarrow X$ is defined by the composite homomorphism $\pi(X, x) \rightarrow G \rightarrow G'$.

Proof. By hypothesis, the bundles $F^{n*}E, n \geq 1$ are all semistable. Since k is a finite field, by boundedness, the set $\{F^{n*}E, n \geq 1\}$ is a finite set. Hence $F^{n*}E \cong F^{(n+\ell)*}E$ for some positive integers n and ℓ . We now consider the monodromy group scheme M_n , and the monodromy bundle E_{M_n} , of the bundle $F^{n*}E$ for $n \geq 0$ (with the convention that $F^{0*}E = E$). Let F_n denote the Frobenius morphism of the group scheme M_n . The isomorphism $F^{n*}E \cong F^{(n+\ell)*}E$ implies that $F_n^\ell(M_n) \cong M_{n+\ell} \cong M_n$, and further the monodromy bundles E_{M_n} and $E_{M_{n+\ell}}$ are isomorphic. We now observe that the isomorphism

$$F_n^\ell(M_n) \cong M_{n+\ell} \cong M_n$$

implies that the group scheme M_n is finite and reduced. Then by Lemma (4.12) in [1], it follows that $M_{0,red}$, the reduced part of the group scheme M_0 (where M_0 denotes the monodromy group scheme of the bundle E), is isomorphic to M_n . Also, if F_0 denotes the Frobenius morphism of M_0 , we obtain $F_0^n(M_0) \cong M_n$. Hence M_0 is a finite group scheme over k . The monodromy bundle E_{M_0} is a principal M_0 bundle on X , and the reduction of structure group $E_{M_0} \subset E$ implies that the pull back of the bundle E to E_{M_0} is trivial. The principal M_0 bundle $E_{M_0} \rightarrow X$ defines a quotient $\pi(X, x) \rightarrow M_0$ (see [5]), and the composite homomorphism

$$\pi(X, x) \rightarrow M_0 \rightarrow G$$

is the homomorphism we want defining the principal G -bundle E .

For the second part, let $G \rightarrow G'$ be a homomorphism of semisimple groups over k . Since G is semisimple, there are three cases to consider:

Case (i) $G \subset G'$ is a closed immersion: in this case, a faithful representation of

G' is also a faithful representation of G , and the result follows from the fact that the monodromy group scheme and the monodromy bundle of E are independent of the choice of representation.

Case (ii) $G \rightarrow G'$ is a central isogeny: we consider the case $G' = G/Z$, where Z is the centre of G . Let M be the monodromy group scheme of E and E_M the monodromy bundle. Let M' be the image of M in G' and $E_{M'}$ the bundle associated to E_M by the projection $M \rightarrow M'$. Then E' has a reduction of structure group $E_{M'} \subset E'$ to M' , of degree zero (see Definition 4.7 in [1]). If H is a group scheme over k and $E_H \subset E'$ a reduction of structure group of E' of degree zero to H , with $H \subset M'$, then taking $\tilde{H}$ to be the inverse image of H in G , we obtain a principal $\tilde{H}$ bundle $E_{\tilde{H}}$ and a reduction of structure group $E_{\tilde{H}} \subset E$ of degree zero with $\tilde{H} \subset M$. This contradicts the minimality of M and E_M (see Corollary (4.10) in [1]). It follows that M' is the minimal subgroup scheme of G' with a degree zero reduction of structure group to M' . Hence, again by Corollary (4.10) in [1], it follows that M' is the monodromy group scheme of E' and $E_{M'}$ is the monodromy bundle of E' .

Case (iii) $G \rightarrow G'$ is purely inseparable: to treat this case, it is enough to consider the Frobenius morphism $F : G \rightarrow G$, as the general case is what is usually called a Frobenius sandwich. Given the principal G bundle $E \rightarrow X$, the bundle induced by the Frobenius $F : G \rightarrow G$ is the pull-back F^*E (where the pullback is taken under the Frobenius morphism of X). Let M be the monodromy group scheme of E and $E_M \subset E$ the

monodromy reduction. We know that M is a finite groupscheme. Let F_M denote the Frobenius morphism of M and let M' be the image of M under F_M . The M bundle E_M induces a M' bundle $E_{M'}$ and a reduction of structure group $E_{M'} \subset F^*E$. If there exists a proper subgroup scheme $N \subset M'$ and a reduction of structure group $E_N \subset E_{M'}$, then the inverse image of E_N under the projection $E_M \rightarrow E_{M'}$ would define a reduction of structure group of E_M , contradicting the minimality of E_M . Therefore $E_{M'}$ is a minimal reduction of structure group of F^*E , and hence is the monodromy reduction.

It now follows that the monodromy of the bundle E' is given by the composite homomorphism $\pi(X, x) \rightarrow G \rightarrow G'$. Q.E.D.

4. When X does not have a rational point

Let X be a smooth projective curve over a finite field k as before and let $E \rightarrow X$ be a principal G bundle on X which is strongly semistable. Let l be a finite Galois extension of k over which X acquires a rational point. Let X_l denote the base change of X to $\text{Spec}(l)$ and E_l the base change of E to $\text{Spec}(l)$. Then by *section 3* above, E_l is defined by a homomorphism $\pi(X_l) \rightarrow G_l$, where $\pi(X_l)$ denotes the fundamental group scheme of X_l and G_l denotes the base change of G to $\text{Spec}(l)$. We have the following exact sequence of group schemes over $\text{Spec}(k)$,

$$1 \rightarrow \pi(X_l, x) \rightarrow \pi(X, x) \rightarrow \text{Gal}(l/k) \rightarrow 1$$

where x is a closed point of X with residue field l , so it is a rational point over l , π is the fundamental group scheme, and $\text{Gal}(l/k)$ is the Galois group

of l over k . Here $\pi(X_l, x)$, which is a group scheme over $\text{Spec}(l)$, is regarded as a group scheme over $\text{Spec}(k)$ by the projection $\text{Spec}(l) \rightarrow \text{Spec}(k)$. The homomorphism $\pi(X_l, x) \rightarrow G_l$ factors through a finite group scheme quotient, say H , of $\pi(X_l, x)$. Since E_l is actually defined over k , it follows that the monodromy group scheme M of E_l , and the monodromy reduction $E_M \subset E_l$, remain invariant under the action of $\text{Gal}(l/k)$ on E_l . Hence the subgroup-scheme $H \subset G_l$ remains invariant under the action of $\text{Gal}(l/k)$. Every group-scheme quotient of $\pi(X_l, x)$ invariant under the action of $\text{Gal}(l/k)$ on $\pi(X_l, x)$ descends to a quotient of $\pi(X, x)$. Hence H is a quotient of $\pi(X, x)$ and we obtain the homomorphism $\pi(X, x) \rightarrow G$ defining E on X over k .

References

1. Indranil Biswas, A.J. Parameswaran and S. Subramanian: Monodromy group for a strongly semistable principal bundle over a curve, *Duke Math. J.* **132**, No.1, 1-48 (2006).
2. H.Esnault, P.H.Hai, and X.Sun, On Nori's fundamental group scheme, arXiv:math.AG/0605645
3. H. Lange and U. Stuhler: Vectorbündel auf Kurven und Darstellungen der algebraischen Fundamentalgruppe, *Math. Z.* **156**, No.1, 73-83 (1977).
4. M.V. Nori: On the representations of the fundamental group scheme, *Comp. Math.*, **33**, 29-41 (1976).

5. M.V. Nori: The fundamental group scheme, *Proc. Ind. Acad. Sci. (Math. Sci.)*, **91**, 73-122 (1982).
6. S.Ramanan and A.Ramanathan: Some remarks on the instability flag, *Tohoku Math. J.(2)*, **36**, 269-291 (1984).
7. J.P. Serre: Galois cohomology, English translation by P.Ion, Springer, 1997

School of Mathematics
Tata Institute of Fundamental Research
Homi Bhabha Road, Colaba
Mumbai 400 005
India
E-mail: subramnn@math.tifr.res.in